



Influence of soil water content and soil amendments on trace metal release and seedling growth in serpentine soil

Viraj Gunarathne¹ · Nishanta Rajakaruna^{2,3} · Udaya Gunarathne¹ · Jayanta Kumar Biswas⁴ · Zach A. Raposo² · Meththika Vithanage¹

Received: 6 February 2019 / Revised: 1 May 2019 / Accepted: 10 May 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Purpose This study was conducted to evaluate the synergistic effects of organic amendments and soil water status on trace metal release from serpentine soil.

Materials and methods Two organic amendments, dendro-biochar (BC) and municipal solid waste compost (CM), were added to serpentine soil at four different ratios, specifically 2.5:0.0, 2.5:1.0, 2.5:2.5, and 2.5:5.0% (w/w). Along with the control (with no organic amendments), each soil treatment was incubated separately under saturated point (SP) and field capacity (FC) water content for 10 days. Subsamples were obtained from each treatment to analyze the bioavailable trace metal concentration and related edaphic parameters, namely total organic carbon (TOC), nitrate (NO_3^-), phosphate (PO_4^{3-}), and cation exchange capacity (CEC). Then, the soil solution was eluted from each treatment and incubated for 10 days under permanent wilting point (PW). Furthermore, a seed germination test was performed under the different treatments.

Results and discussion Significant reductions ($p < 0.05$) in bioavailable concentration of all four trace metals were observed in all the amendment ratios under all water status treatments (SP, FC, SP-PW, and FC-PW), compared with the control. Furthermore, FC-PW with the highest amendment ratio (2.5% BC:5.0% CM) reduced Ni by 67.6%; FC-PW with 2.5% BC + 2.5% CM immobilized Mn and Co by 92.1 and 96.9%, respectively, and SP water status with all four amendment ratios immobilized 100% of bioavailable Cr. Maximum amendment ratio under all four water status enhanced %TOC and significantly increased PO_4^{3-} concentration in SP-PW. However, FC showed comparatively high NO_3^- concentration than other treatments. Germination index (GI) for mung beans and tomato did not show a significant difference in response to amendment ratios or soil water status; however, SP treatment expressed significantly high seedling vigor (SVI) for mung beans.

Conclusions Treatments of BC and CM effectively immobilize the bioavailable fraction of trace metals in serpentine soils. Increasing amendment ratio increases the %TOC regardless of the soil water status, whereas SP-PW is favorable for the availability of PO_4^{3-} , and FC is favorable for availability of NO_3^- . The GI for mung beans and tomato seed was not influenced by the soil water status nor by the amendment ratios. However, the SVI of mung bean seedlings was controlled by the soil water status.

Keywords Biochar · Compost · Serpentine · Plant nutrients · Remediation · Soil water content · Trace metals

Responsible editor: Kitae Baek

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11368-019-02349-9>) contains supplementary material, which is available to authorized users.

✉ Meththika Vithanage
meththika@sjp.ac.lk

¹ Ecosphere Resilience Research Centre, Faculty of Applied Sciences, University of Sri Jayewardenepura, Nugegoda, Sri Lanka

² Biological Sciences Department, California Polytechnic State University, San Luis Obispo, CA 93407, USA

³ Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa

⁴ Enviromicrobiology, Ecotoxicology and Ecotechnology Research Laboratory, Department of Ecological Studies and International Centre for Ecological Engineering, University of Kalyani, Kalyani, West Bengal 741235, India

1 Introduction

Ultramafic rocks and related serpentine soils are distributed all over the world and represent nearly 1% of surface area of the Earth (Vithanage et al. 2014). Even though these soils show sparse distribution, their occurrence is reported in all major biomes and all the continents except Antarctica (Harrison and Kruckeberg 2008). Natural weathering processes of ultramafic rocks lead to pedogenesis of serpentine soils and sediments characterized by a high metal content (i.e., Mg, Ni, Cr, Co, and Mn), high Mg:Ca ratios, and nutrient poor conditions (Rajapaksha et al. 2012; Vithanage et al. 2014; Kumarathilaka et al. 2016). Rajapaksha et al. (2012) found 6179, 1036, 11,352, and 121 mg kg⁻¹ respective values for total concentration of Ni, Mn, Cr, and Co in < 2-mm fraction of soil from Ussangoda serpentine outcrop, Sri Lanka. Some of the industrialized and most populated human settlements in North America, Europe, Africa, Southeast Asia, and Australia are on or in close proximity to extensive serpentine outcrops (Brooks 1987; Harrison and Rajakaruna 2011; Kumarathilaka et al. 2016). In addition, some serpentine deposits have been extended into agricultural regions (Ye-Tao et al. 2012; Neilson and Rajakaruna 2015), including in Mouriki-Thiva, Central Greece, where Antibachi et al. (2012) found 621–2639 and 134–856 mg kg⁻¹ of Ni and Cr in soil, respectively. Moreover, due to high rainfall and erosion, serpentine soil and sediments can separate from its main body, transport through irrigation systems, and deposit in valleys and lowlands which are used extensively for agriculture (Wei et al. 2018). Ranau Valley in Malaysia was reported to have Mn (43,063.8 mg kg⁻¹), Cr (7332.2 mg kg⁻¹), Ni (6322.3 mg kg⁻¹), and Co (1004.7 mg kg⁻¹) in paddy soil due to erosion, transport, and sedimentation (Aziz et al. 2015). Similarly, the serpentine sediment deposition in agricultural areas located in floodplains of Whatcom County, Washington, USA, had resulted from a natural landslide associated with headwater areas of the Swift Creek tributary (Baugé et al. 2013).

Due to the presence of elevated levels of trace metals, serpentine soils and sediments can be considered as non-anthropogenic sources of trace metal contamination (Rajakaruna and Baker 2004; Seneviratne et al. 2015). Many studies have found extreme total concentrations of trace metals, especially Ni, Cr, Mn, and Co in serpentine soils in various localities worldwide (Lázaro et al. 2006; Antibachi et al. 2012; Aziz et al. 2015; Galey et al. 2017). However, the total trace metal concentration does not accurately reflect the extent of environmental pollution and ecological risks because trace metals show fractionation among different components of soil (Cheng et al. 2011; Tashakor et al. 2011). Even though the soil is divided into different fractions, including (1) adsorptive and exchangeable, (2) contained in carbonate phase, (3) bound to reducible phases (Fe and Mn oxides),

(4) occluded with organic matter and sulfides, and (5) residual/lattice metals, and the trace metals are distributed among these fractions (Kierczak et al. 2008), the bioavailability, mobility, and toxicity are solely dependent on the amount of trace metal bound in the adsorptive and exchangeable fraction (Kabala and Singh 2001). The study done by Rajapaksha et al. (2012) found 167.6 (± 6.4) and 45.5 (± 3.0) mg kg⁻¹, respectively, of bioavailable Ni and Mg from serpentine sediments. Further, the same study revealed the respective rates of 1.55×10^{-13} and 7.89×10^{-14} mol m⁻² s⁻¹ of Ni and Mn release into water within 24 h.

Soil moisture regimes have a direct effect on many physicochemical and microbiological qualities of soils, including organic matter content, pH, and electrical conductivity (Van den Berg and Loch 2000; Angle et al. 2003; Gray and McLaren 2006; Zheng and Zhang 2011) and are indirectly involved in the fractionation of metals and their bioavailability. Zheng and Zhang (2011) demonstrated the effect of hydrological regimes such as field capacity, flooding, and wetting-drying cycles for transformation rates of trace metals among different fractions of paddy soils. The flooding condition of soil contributes to higher transformation rates of trace metals towards stable fractions (i.e., Fe–Mn oxide and organic matter bound) than soil incubated under 75% of field capacity and soils which are kept under wetting–drying cycles. Trace metal transformation into stable fractions under flooded conditions is governed by increased pH levels, metal ion precipitation with sulfides, and increased concentration of iron oxides which are induced by the saturated water level (Zheng and Zhang 2011). Therefore, current soil water status such as field capacity and wilting point as well as the dynamics of soil moisture content should be taken into account in order to assess the release and bioavailability of trace metals from certain soil types.

Organic amendments have been utilized as fertilizer and considered as a cost-effective approach to immobilize the bioavailable trace metal fraction and mitigate the negative consequences associated with trace metals in contaminated soils (Almaroai et al. 2014; Herath et al. 2015; Rizwan et al. 2017). In this regard, compost and biochar are known to be agents with high potential to adsorb trace metals (Almaroai et al. 2014). Compost is a biologically stable organic compound produced from biological oxidation and humification of raw organic matter which are mostly the waste products from agriculture or food industry (Bernal et al. 2007). Many studies have recognized the capacity of compost as an amendment for remediation of trace metal contaminated sites (Kargar et al. 2015; Liang et al. 2017). It is also considered as a nutrient rich organic fertilizer that can be effectively used to improve soil quality and microbial activities (Lamey and Angers 2012; Mackie et al. 2015; Cerdeira-Pérez et al. 2019).

Characteristics of serpentine soil are distinct (e.g., high Mg:Ca ratio) from any other soils anthropogenically contaminated with trace metals (Rajakaruna and Boyd 2014).

Although the soil water status influences bioavailability of metals, not many studies have focused on the influence of soil moisture on trace metal availability in serpentine soils (Angle et al. 2003; Aziz et al. 2015; Tashakor et al. 2017). This is particularly significant as serpentine soils are being increasingly used for cultivation (Fernandez et al. 1999; Kanellopoulos et al. 2015; Nkrumah et al. 2019; Cerdeira-Pérez et al. 2019). As the incorporation of organic amendments under different soil moisture regimes may induce differential trace metal release, this study investigated the influence of soil water status on the bioavailability of trace metals in serpentine soil and evaluated the effects of different bioamendments on immobilization/release of trace metals, including on the germination and vigor of seedlings.

2 Materials and methods

2.1 Study site and sample collection

Serpentine soil used for the study was obtained from Ussangoda serpentine outcrop (latitude 6° 05' 54" N and longitude 80° 59' 11" E) located in the Southern part of Sri Lanka (see Fig. 1 in Vithanage et al. (2014)). Soil sample collection was done at several random locations and mixed together to acquire a homogenized composite sample weighing approximately 20 kg. This composite soil sample was used for all experiments.

2.2 Incubation experiment

Soil treatments consisted of the following: (1) control (serpentine soil only), (2) soil + 2.5% dendro-biochar (*Gliricidia sepium*) (BC), (3) soil + 2.5% BC + 1.0% municipal solid waste compost (CM), (4) soil + 2.5% BC + 2.5% CM, and (5) soil + 2.5% BC + 5.0% CM. The samples within each treatment were incubated under two different soil water status; saturated point (SP) and field capacity (FC) to evaluate the influence of soil water content and different amendments/amendment ratios on the metal release from serpentine soils. All incubation tests were done in plastic containers, and each treatment was maintained in triplicate. For each treatment, 450 g of serpentine soils was used and compacted to achieve 1.48 g cm^{-3} of bulk density. Consequently, for each treatment, deionized water was added until SP. Containers having SP samples were air tightened by covering them with polyethylene and parafilm. The FC samples in porous containers were allowed to drain excess water for 3 days until no more water leached out, thus, reaching field capacity. Afterwards, containers having FC samples were also air tightened as mentioned earlier for SP samples. After 10 days of incubation, subsamples were taken from each soil sample to perform further experiments. Soil solution of each treatment was then

eluded under 1 atm of pressure. A fraction of the subsamples from each treatment pot was separated for further chemical analysis. Subsequently, after pore water extraction, the incubation of remaining SP and FC soil samples was extended for 10 more days at permanent wilting point (PW), and after completion of incubation, subsamples were taken to carry out further experiments.

2.3 Diethylenetriaminepentaacetic acid extraction

The extraction of the bioavailable fraction of trace metals was performed using diethylenetriaminepentaacetic acid (DTPA) extraction method as described in Lindsay and Norvell (1978). Twenty milliliters of 0.005 M DTPA, 0.01 M CaCl_2 , and 0.1 M triethanolamine solution was poured into 10 g of each air-dried subsample of SP, FC, and PW treatments and shaken for 2 h at 100 rpm. Resulting solutions were centrifuged at 2000 rpm for 15 min and filtered through 0.45 μm filter paper. Filtrates were analyzed for Ni, Mn, Co, and Cr using atomic absorption spectroscopy (AAS) (Shimadzu AA-7000; Japan).

2.4 Ammonium acetate extraction

Cation exchange capacity (CEC) of each treatment was evaluated using the ammonium acetate extraction procedure (Anderson and Ingram 1993). Two grams of air-dried soil samples from each treatment was transferred into separate plastic tubes, and 20 ml of ammonium acetate solution was added. After shaking at 100 rpm for 15 min, centrifugation was performed at 2000 rpm for 15 min, and supernatants were filtered through 0.45- μm syringe filters. Filtrate was analyzed by AAS for Na, K, Ca, and Mg, and CEC of each treatment was calculated using the obtained data.

2.5 Total organic carbon, soil available phosphate, and nitrate

Nutrient contents (i.e., total organic carbon, available P, and N) of each SP, FC, and PW treatment we evaluated using the standard colorimetric methods described in Anderson and Ingram (1993).

2.6 Leachate analysis

Leachates that eluded from each treatment of SP and FC soils were filtered through 0.45 μm syringe filters and analyzed using AAS for Ni, Mn, Co, and Cr.

2.7 Seed germination experiment

Seed germination study was carried out to evaluate the synergistic effect of soil water status and different amendments/

amendment ratios on germination performance of two mostly cultivated species in Sri Lanka, namely, Mung bean (*Vigna radiata*) and Tomato (*Solanum lycopersicum*). Twenty grams of serpentine soils in separate petri dishes was incubated for 7 days with five different amendment ratios used in the previous study (i.e., (1) control (serpentine soil only), (2) soil + 2.5% BC, (3) soil + 2.5% BC + 1.0% CM, (4) soil + 2.5% BC + 2.5% CM, and (5) soil + 2.5% BC + 5.0% CM) with respective soil water status of SP, FC, and PW. Each treatment was run as triplicates. After the incubation period, each petri dish was divided into equal parts, and five seeds each of mung bean and tomato were placed separately in each area. Seeds were allowed to germinate under ambient atmospheric conditions. The germination percentage of seeds in each setup was recorded at 24 h intervals for 5 days. Seeds were considered as germinated when the radicle emerged. After 5 days of germination, the length of each seedling was measured, and the germination index, the time for 50% germination, and the seedling vigor index were calculated.

2.7.1 Germination index

Germinations index (GI) for each seed type was calculated using the equation below (Siddiqui and Al-Wahaibi 2014).

$$GI = \sum Gt/Dt \quad (1)$$

where Gt is germination percentage of seeds and Dt is number of days taken by seeds to germinate.

2.7.2 Seedling vigor index

Seedling vigor index (SVI) was calculated according to Vashisth and Nagarajan (2010).

$$SVI = \text{germination percentage} \times \text{mean value of seedling length} \quad (2)$$

2.8 Statistical analysis

All statistical analyses were carried out using the statistical software package MINITAB® 16.1.0. Bioavailable trace metals (viz. Ni, Mn, Co, and Cr), soil nutrients (viz. NO_3^- , PO_4^{3-} , and %TOC), CEC, seed germination index, and seedling vigor index among five different amendment ratios under different soil water status were compared performing one-way ANOVA. Additionally, for each measured variable, Tukey pairwise comparisons were performed among five amendment ratios.

3 Results

Table 1 highlights the DTPA extractable trace metal concentration of Ussangoda serpentine soils. The respective percentage values for bioavailable Ni, Mn, Cr, and Co of the total concentrations of metals mentioned in the previous study of Rajapaksha et al. (2012) are 2.007, 7.581, 0.005, and 5.900%. The serpentine soil examined contained comparatively low concentrations of NO_3^- and PO_4^{3-} which are considered as available forms of nitrogen and phosphorus for the plants. Similarly, low values for percentage total organic carbon (%TOC) and cation exchange capacity (CEC) were reported for the serpentine soils (Table 1).

Figure 1 illustrates the influence of soil water content on bioavailability of trace metals: Ni, Mn, Cr, and Co. The highest bioavailable concentrations of Ni and Mn (133.72 and 0.95 mg kg^{-1} , respectively) were reported in SP-PW. However, the lowest concentrations were observed at FC for Ni ($121.39 \text{ mg kg}^{-1}$) and SP for Cr (0.20 mg kg^{-1}). In contrast, the highest bioavailability of Mn and Co was found at SP (71.40 and 5.90 mg kg^{-1} , respectively) and the lowest bioavailable concentration was reported at SP-PW (45.91 and 4.05 mg kg^{-1} , respectively).

Application of organic amendment at all four ratios significantly reduced (one-way ANOVA, $p < 0.0001$ at $\alpha = 0.05$) the bioavailable concentration of all four examined trace metals under all four soil water contents (Fig. 2). The maximum bioavailable concentration of Ni was observed at SP-PW with no amendment (control) ($133.72 \text{ mg kg}^{-1}$) and the lowest was present at FC-PW with the highest amendment ratio (43.32 mg kg^{-1}). The reduction of bioavailable Ni concentration in FC-PW soil at the maximum amendment ratio is 67.6% compared with its control. However, considering the overall results, 2.5% BC + 2.5% CM and 2.5% BC + 5.0% CM were equally effective for immobilization of bioavailable Ni from serpentine soil.

The control set at SP contained significantly higher (one-way ANOVA, $p < 0.050$ at $\alpha = 0.05$) bioavailable concentration of Mn (71.40 mg kg^{-1}) (Fig. 2). The minimum value for bioavailable Mn concentration was reported from FC-PW treated with 2.5% BC + 1.0% CM amendment ratio (5.64 mg kg^{-1}), which is a 92.1% reduction from the control treatment of FC-PW. However, statistical analyses revealed that the SP-PW and FC-PW with all amendment ratios (i.e., 2.5% BC + 0.0–5.0% CM) exhibited no significant difference (one-way ANOVA, $p > 0.050$ at $\alpha = 0.05$) and all were found equally effective for immobilization of Mn in the serpentine soil.

The highest DTPA extractable Co concentration was reported in the control set (with no amendment) at SP (5.88 mg kg^{-1}). However, it was not significantly different (one-way ANOVA, $p > 0.050$ at $\alpha = 0.05$) from the control at FC (5.42 mg kg^{-1}). Similar to Mn, the lowest concentration

Table 1 Elemental composition and physicochemical characteristics of serpentine soil (< 2 mm) and the organic amendments

DTPA extractable metal concentration (mg/kg)				Ammonium acetate extractable cation concentration (mg/kg)			pH	NO ₃ ⁻ (mg kg ⁻¹)	PO ₄ ³⁻ (mg kg ⁻¹)	TOC (%)	CEC (cmol _c kg ⁻¹)	Soil moisture content (%)	Field capacity	
Ni	Mn	Cr	Co	Na	K	Ca	Mg							
124.00 ± 9.62	78.54 ± 4.21	0.60 ± 0.02	7.14 ± 0.14	1.85 ± 0.04	191.05 ± 4.91	18.16 ± 2.26	22.19 ± 5.12	6.68 ± 0.14	8.75 ± 0.63	9.12 ± 0.94	2.00 ± 1.09	1.12 ± 0.08	34.40	26.47
DTPA extractable metal concentration (mgkg ⁻¹)														
				pH			EC (μS cm ⁻¹)	NO ₃ ⁻ (mgkg ⁻¹)	PO ₄ ³⁻ (mgkg ⁻¹)	TOC (%)	CEC (cmol _c kg ⁻¹)			
Ni	Mn	Cr	Co											
Dendro-biochar	N/A	N/A	N/A	N/A	10.29 ± 0.21		1488 ± 4.50	1.80 ± 0.51	71.70 ± 1.51	5.92 ± 0.94	52.14 ± 1.02			
MSW compost	0.03 ± 0.00	38.88 ± 3.81	ND	0.67 ± 0.08	8.00 ± 0.12		2620 ± 6.30	3.90 ± 0.47	8.78 ± 0.91	38.52 ± 1.04	49.08 ± 1.06			

Value proceeding ± is the standard deviation (SD) for that element or method

of bioavailable Co was observed from the treatment at FC-PW amended with 2.5% BC + 1.0% CM (0.18 mg kg⁻¹), a 96.9% reduction from the control set at FC-PW.

As shown in Fig. 2, Cr exhibits the lowest bioavailability compared with the other three trace metals. The highest bioavailability of Cr was found at SP-PW with no amendments (0.95 mg kg⁻¹). However, it is not significantly different (one-way ANOVA, $p > 0.050$ at $\alpha = 0.05$) from the control at FC-PW. Interestingly, the SP soils incubated with any amendment ratio (i.e., 2.5% BC + 0.0–5.0% CM) were able to completely immobilize Cr.

The soil solution eluted from each SP and FC treatments after 10 days of incubation contained extremely low concentrations of Ni and Mn compared with their bioavailable concentrations (Table 2). Moreover, the Ni concentration of both SP and FC solutions decreased with the addition of amendments. The FC soils showed continuous reduction of Ni concentration until FC 4 (2.5% BC + 2.5% CM) and became constant afterwards. However, the lowest Ni concentration was reported in SP 3 (2.5% BC + 1.0% CM) and increased slightly with the increasing amendment ratio. In contrast to Ni, soil solution from SP 1 (control) contained lowest Mn concentration, and the concentration increased with an increasing amendment ratio. Besides, in FC soils, the highest concentration of Mn was observed in FC 1 (control) while minimum concentration was found in FC 3 (2.5% BC + 1.0% CM) and increased subsequently. Moreover, the Mn concentrations in soil solutions from FC soils were lower than those in extracted soil solution from SP soils, except SP 1. Furthermore, the Co and Cr were not present in either of the extracted soil solutions.

With the addition of organic amendments, dendro-biochar (BC) and municipal solid waste compost (CM), soil nutrient content and examined soil chemical parameters increased drastically (Fig. 3). The highest amendment ratio resulted in the highest (one-way ANOVA, $p < 0.050$ at $\alpha = 0.05$) nutrient content of treated serpentine soil except FC and SP-PW treatments which did not exhibit significant difference among amendment ratios of 2.5% BC, 2.5% BC + 2.5% CM, and 2.5% BC + 5.0% CM (one-way ANOVA, $p > 0.05$ at $\alpha = 0.05$). However, the maximum amendment ratio in FC-PW soil contained the highest concentration (32.49 mg kg⁻¹) of NO₃⁻ among all the treatments, and it was 271.3% higher than in the untreated serpentine soil. In contrast to the NO₃⁻, the highest PO₄³⁻ concentration was observed in SP-PW soil with the uppermost amendment ratio (15.48 mg kg⁻¹), which is 69.7% higher compared with untreated serpentine soils. Moreover, the overall results show that the conditions provided by SP-PW water status are more favorable for high PO₄³⁻ concentration than the other three soil water status (Fig. 3).

As expected, %TOC of treated soils increased with the increasing amendment ratio (Fig. 3). However, FC-PW with maximum amendment ratio contained the highest amount of




Fig. 1 Concentrations of bioavailable trace metals of Ni, Mn, Co, and Cr in untreated serpentine soil under different soil water contents. Letters are assigned from the pairwise comparison performed using Tukey method.

%TOC (3.2%), which is 58.0% higher than in the untreated serpentine soils. Moreover, some of the treatments such as 2.5% BC + 2.5% CM amendment ratios of SP-PW and FC-PW soil water status contributed to reduced %TOC than that of their consecutively low amendment ratio.

Furthermore, all of the treated serpentine soil samples with organic amendments clearly showed increased CEC values compared with the untreated serpentine soil (Fig. 3). The highest CEC value was reported for SP soil with the highest amendment ratio ($8.29 \text{ cmol}_c \text{ kg}^{-1}$) while the lowest CEC value was observed in FC treatment with no amendment ($1.65 \text{ cmol}_c \text{ kg}^{-1}$). For instance, the highest CEC value of SP treatment indicated 640.5% of increment over the untreated serpentine soil.

The mung beans and tomato seeds germinated in both SP and FC soil while no seed germination was observed in the soil incubated at WP (Table S1 - Electronic Supplementary Material). All the mung beans in soils incubated at FC soil water status germinated regardless of the amendment ratio. Therefore, the germination index (GI) for mung seed that

(SP, incubated at saturated point; FC, incubated at field capacity; SP-PW, incubated at saturated point followed by permanent wilting point; and FC-PW, incubated at field capacity followed by permanent wilting point)

germinated at FC exhibited no variation since all values are identical (Fig. 4). Also, the GI among mung beans germinated with different amendment ratios under SP soil water status did not show any significant difference (one-way ANOVA, $p = 0.452$ at $\alpha = 0.05$). Although, the germination of tomato seeds showed differences between SP and FC soil water status and among amendment ratios, there was no significant difference (one-way ANOVA, $p > 0.050$ at $\alpha = 0.05$) for GI (Fig. 4).

Figure 5 shows the variation of seedling vigor index (SVI) among mung bean and tomato seedlings which were grown in SP and FC soils with different amendment ratios. Although GI did not reflect a significant difference for the germination of mung bean between SP and FC treatments, SVI indicated a significant difference (one-way ANOVA, $p < 0.0001$ at $\alpha = 0.05$) for the vigor of mung bean seedlings among SP and FC. Conversely, different amendment ratios within SP and FC treatments did not show significant effect on the vigor of mung bean seedlings (one-way ANOVA, $p = 0.430$ and $p = 0.111$ at $\alpha = 0.05$ for SP and FC, respectively). In contrast,




Fig. 2 Bioavailable trace metal concentrations at different amendment ratios under different soil water status. Letters are assigned from the pairwise comparison performed for five treatments under different soil water levels using Tukey method. (SP, incubated at saturated point; FC,

incubated at field capacity; SP-PW, incubated at saturated point followed by permanent wilting point; and FC-PW, incubated at field capacity followed by permanent wilting point)

SVI of tomato reflected no difference for SP and FC treatments (one-way ANOVA, $p = 0.258$ at $\alpha = 0.05$). Furthermore, different amendment ratios of SP had significant influence (one-way ANOVA, $p = 0.020$ at $\alpha = 0.05$) on the SVI of tomato while different ratios of amendment used for

FC did not exhibit a significant difference (one-way ANOVA, $p = 0.063$ at $\alpha = 0.05$) on the vigor of tomato seedlings. Therefore, water status of serpentine soil is an important determinant for the vigor of mung bean seedlings on serpentine soils, however, not for that of tomato seedlings.

Table 2 Trace metal concentration in soil solution eluted from SP and FC soils

Soil solution		SP1	SP2	SP3	SP4	SP5	FC1	FC2	FC3	FC4	FC5
Metal concentration (mg kg ⁻¹)	Ni	0.87	0.23	0.21	0.26	0.32	0.94	0.59	0.41	0.28	0.28
		±0.03	±0.02	±0.02	±0.01	±0.05	±0.03	±0.01	±0.04	±0.01	±0.01
	Mn	0.08	0.59	0.67	1.12	0.88	0.94	0.34	0.25	0.41	0.40
		±0.01	±0.01	±0.02	±0.04	±0.01	±0.02	±0.02	±0.01	±0.01	±0.03

SP and FC indicate two soil water status saturated point and field capacity while no. 1–5 represent ratio of organic amendment

Values indicated are average of three determinations and the values proceeding ± is the standard deviation (SD)

*Co and Cr are not detected in any of the soil solutions eluted




Fig. 3 Changes in the concentration of nitrate (NO₃⁻), phosphate (PO₄³⁻), total organic carbon (%TOC), and cation exchange capacity (CEC) at different amendments ratios under different soil water status. Letters are assigned from the pairwise comparison performed for five treatments under

different soil water levels using Tukey method. (SP, incubated at saturated point; FC, incubated at field capacity; SP-PW, incubated at saturated point followed by permanent wilting point; and FC-PW, incubated at field capacity followed by permanent wilting point)

4 Discussion

The bioavailable concentrations found for Ni, Mn, Cr, and Co (Table 1) are comparatively lower when compared with the total trace metal concentrations reported by Rajapaksha et al. (2012) for the soils in the same geological region. Metal bioavailability is the factor that exclusively determines the magnitude of the risk generated by toxic heavy metals for the environment and for humans (van Gestel 2008). Therefore, the bioavailable metal concentration indicates more accurate information about environmental toxicity generated by toxic trace metals. As a fraction of the total metal content, bioavailable trace metal concentration is lower than total trace metal concentration of soil. The DTPA extraction procedure delivers reliable estimation of bioavailable trace metal concentration in soils (Peijnenburg et al. 2007). The bioavailable Ni, Mn, Cr, and Co in the DTPA extraction of Ussangoda serpentine soil

were 2.007, 7.581, 0.005, and 5.900% of the total metal concentration, respectively. The findings of Kumarathilaka and Vithanage (2017) reflect a similar tendency as bioavailable Ni and Mn concentrations of serpentine soils were 0.30 and 0.25%, respectively, of total concentrations. Moreover, 33–323 and 11–76 mg kg⁻¹ of Ni and Mn, respectively, were found as bioavailable fractions from four serpentine localities in Sri Lanka (Vithanage et al. 2014). Even though the bioavailable concentration is comparatively lower than the total trace metal concentration, long-term risks associated with their accumulation in food chains cannot be ignored.

Serpentine soils are often characterized as nutrient poor, especially with low amounts of N, P, K, and Ca (Kruckeberg 2002; Vithanage et al. 2014; Baumeister et al. 2015). However, the low N conditions are mostly applicable only to the serpentine localities with scarce vegetation (Alexander et al. 2007), such as the Ussangoda serpentine body




Fig. 4 Effects of soil water content and amendment ratio on germination index (GI) of mung bean and tomato. Letters are assigned from the pairwise comparison performed for five treatments under different soil

water levels using Tukey method. (SP, incubated at saturated point and FC, incubated at field capacity)

(Rajakaruna and Bohm 2002). Even though serpentine soils have elevated Mg levels, amounts of Ca, Na, and K are exceptionally low (Kumar and Maiti 2013). Therefore, low Ca, Na, and K concentrations ultimately contribute to low CEC values. Serpentine soils usually contain low organic matter content (Proctor and Woodell 1975; Seneviratne et al. 2015), typically attributed to poor

vegetation cover. Seneviratne et al. (2015) found 2.3% total organic carbon in soil from Yudhaganawa serpentine deposit in Sri Lanka, which is similar to the findings of the present study. Moreover, the comparatively low soil water holding capacities observed at saturation point and field capacity are often reflected in serpentine soils worldwide (Schedlbauer 2015).




Fig. 5 Effects of soil water content and amendment ratio on seedling vigor index (SVI) on serpentine soils. Letters are assigned from the pairwise comparison performed for five treatments under different soil

water levels using Tukey method. (SP, incubated at saturated point and FC, incubated at field capacity)

Soil moisture regime directly causes for changes in physicochemical and biological attributes of soil, including pH, electrical conductivity, and organic matter content (Van den Berg and Loch 2000). Kashem and Singh (2004) list a number of changes which occur in soil under flooded conditions, including a decrease in redox potential (Eh) due to microbial respiration, increase in soil pH, expansion of oxide surfaces, decrease in crystallinity, and the occurrence of hydrated forms of Mn and Fe. Changes of such physicochemical and biological factors influence the redistribution of trace metals among different fractions of soil as well as the dissolution of trace metals (Du Laing et al. 2007; Zheng and Zhang 2011; Liang et al. 2017). In the present study, the soil solutions eluted from serpentine soils incubated at saturated point (SP) and field capacity (FC) for 10 days contained lower concentrations of Ni and Mn compared with their bioavailable concentrations. Co and Cr were not detected in any of the extracted soil solutions. It is hypothesized that Ni is more liable to release in oxygen-rich conditions associated with dissolution of calcium carbonate (Du Laing et al. 2007). But our results do not reflect a pronounced increment in Ni concentration in water extracted from FC incubation compared with SP. In order to detect substantially high Ni concentrations in pore water of FC incubation, it may be necessary to have a considerably long period of time, as the dissolution of calcium carbonate is a slow process governed by weak acidity. Gambrell (1994) suggested that submerged conditions increase the Mn concentration in pore water as a result of microbial reduction of manganese oxides. Our study indicated contradictory results with slightly high concentration of Mn in FC soil solution than that of SP. However, metal dissolution under a reduced environment generated by microbial respiration and subsequent distribution into more stable fractions of soil is a two way process (Han and Banin 1997; Kashem and Singh 2004). Therefore, released Mn under saturated conditions can transform into more stable fractions of soil, leading to a reduction of Mn concentration in the soil solution. Moreover, the comparatively high concentration of Mn (38.88 mg kg^{-1}) present in municipal solid waste compost (CM) might have contributed to the increase of Mn concentration in soil solution with increasing amendment ratio, by dissolving under excess water content present in the saturated point (SP) soils. This is due to the high water content in SP soils which facilitates relatively high dissolution of Mn from soil and CM amendments. However, the bioavailable concentrations of Co and Cr content were comparatively much lower in serpentine soils (Table 1) and consequently were not detected in either SP or FC soil solutions.

The present study indicates the reduction of Mn and Co bioavailability under permanent wilting point (PW) incubations of SP-PW and FC-PW than those of saturated point (SP) and field capacity (FC) incubations (Fig. 1). Under saturated soil conditions, trace metal release is induced by the reduction of Fe and Mn oxides (Han and Banin 2000; Han

et al. 2001). The reduction in bioavailable Mn and Co levels in SP-PW and FC-PW can be attributed to the inhibited microbial reduction by water deficient conditions under permanent wilting point and oxidation under oxic conditions. Conversely, Cr shows declined bioavailability in SP and FC incubations than subsequent incubations at permanent wilting point (SP-PW and FC-PW). Under low redox (reduced) conditions with pH ranging from 2 to 14, Cr forms more stable species with less bioavailability (Avudainayagam et al. 2003). However, soil moisture content has less influence on the redistribution of Cr among different fractions of soil (Han et al. 2003). The bioavailability of Ni showed a significant reduction (one-way ANOVA, $p < 0.0001$ at $\alpha = 0.05$) in FC incubation than in the other three incubations. However, the bioavailable Ni concentrations in all four incubations ranged from 120 to 135 mg kg^{-1} which is closer to the bioavailable Ni concentration in un-incubated serpentine soil ($124.04 \text{ mg kg}^{-1}$, Table 1). The results of Fengxiang and Banin (1997) indicate that the transformation of Ni occurs from exchangeable and carbonate bound fractions to organic matter bound and easily reducible oxide fractions under flooded conditions. Similarly, field capacity conditions are also involved in the redistribution of Ni from exchangeable and carbonate bound fractions to an easily reducible fraction and organic matter bound fractions in a slow manner. Therefore, saturated conditions lead to a more pronounced distribution of Ni into more stable fractions than at field capacity. Similarly, Kashem and Singh (2004) found 47% reduction of ammonium acetate (1 M NH_4OAc , pH 7) extractable Ni in exchangeable fraction after submergence of contaminated soil for 24 weeks. However, 10 days of incubation might not be long enough for the initiation of transformation.

Biochar, a type of black carbon resulting from pyrolysis of biomass under no/limited oxygen conditions, has been proven to be an effective and low-cost organic amendment for trace metal-contaminated sites (Inyang et al. 2016). Previous research of our group has revealed strong immobilization capacity of 2.5% dendro-biochar (*Gliricidia sepium* biochar) on trace metals released from serpentine soils (Herath et al. 2015; Kumarathilaka and Vithanage 2017). Herath et al. (2015) found that dendro-biochar reduces bioavailable Cr, Ni, and Mn in serpentine soil by 96, 36, and 21%, respectively. Similarly, compost generated from various feedstocks undergoing microbial oxidation under anaerobic conditions, is also a suitable amendment for trace metal-contaminated sites (Beesley et al. 2010; Karami et al. 2011). Compost amendment into contaminated soils can reduce bioavailability of trace metals by transformation to a more stable organic matter bound fraction (Manios et al. 2003; O'Dell et al. 2007). Therefore, our study focused primarily on the synergistic effects of different ratios of dendro-biochar (BC) and municipal solid waste compost (CM) with different soil water regimes

for determining the bioavailability of trace metals from serpentine soil.

In our study, all the amendment ratios, ranging from BC:CM ratios of 2.5:0–5%, were involved in a significant reduction (one-way ANOVA, $p < 0.0001$ at $\alpha = 0.05$) of the bioavailability of all four metals examined. Extensive surface area associated with microporous structure, polyaromatic nature, diverse surface functional groups, and high cation exchange capacity make biochar a good adsorbent for trace metals (Tang et al. 2012; Xu et al. 2014). Biochar application immobilizes and increases the stability of trace metals by complexation with surface functional groups, specific and non-specific adsorption, and ion exchange and precipitation (Zhang et al. 2013). In the current study, biochar alone leads to increased immobilization of Co under SP-PW and FC-PW (Fig. 2). However, in most of the treatments, BC and CM together contribute to the maximum immobilization of bioavailable trace metals (Fig. 2). Liang et al. (2017) provide similar results to those of our study, showing that biochar-compost composites immobilize trace metals significantly more than the control. Combination of organic amendments is highly effective in reducing mobility of exchangeable trace metals in soils (Beesley et al. 2014). The humic substances and microorganisms in compost contribute to further oxidation of biochar surfaces, while biochar can also induce the humification of compost, providing favorable conditions for microbial growth (Wu et al. 2016; Wu et al. 2017). Organic matter also has the ability to interact with metals and contributes to their speciation, thereby reducing their mobility. Thus, high organic matter and microbial content in compost can induce high stability and low mobility of trace metals (Kalbitz and Wennrich 1998; Clemente et al. 2006). Further, the comparatively high pH of dendro-biochar (pH 10.29) and municipal solid waste compost (pH 8.00) (Table 1) can synergistically increase the pH of serpentine soil which can then contribute to the stabilization of trace metals. Solubility of trace metals is strongly correlated with soil pH, with high solubility and mobility occurring under low pH values (Huang et al. 2014; Zeng et al. 2011).

All the organic amendment ratios used under all the soil water status significantly (one-way ANOVA, $p < 0.0001$ at $\alpha = 0.05$) increased the concentration of soil NO_3^- , PO_4^{3-} , percent total organic carbon (%TOC), and cation exchange capacity (CEC), compared with the control (Fig. 3). The application of biochar is known to improve various soil properties such as cation exchange capacity (CEC), soil total organic carbon (%TOC), pH, and soil nutrient concentrations (Agegnehu et al. 2015; Zeng et al. 2015). However, the effects of biochar on soil nutrient content are dependent on the type of biomass used for biochar production (Gunaratne et al. 2017). Similarly, compost is a quality organic fertilizer which can provide soil nutrients as well as improve physicochemical and microbial properties of soil (Schulz et al. 2013; Mackie et al. 2015). Figure 3 shows a similar trend for CEC and %TOC, as they increase with increasing amendment ratios and are not influenced by the soil water status.

The dendro-biochar and municipal solid waste compost used for this study consisted of 46.6- and 43.8-fold higher cation exchange capacity (CEC), respectively, than the CEC of serpentine soils (Table 1). The higher CEC values correspond to a higher number of cation exchange sites, indicating the capacity of the amendment to immobilize trace metals by ion exchange. Figure 3 shows the extensive increment of soil total organic carbon (%TOC) levels with an increasing amendment ratio. The dendro-biochar and municipal solid waste compost consist of 2.9- and 19.3-fold higher %TOC, respectively, compared with the serpentine soils (Table 1), contributing to an increase in %TOC in soil with an increasing organic amendment ratio. Increasing organic matter content has a pronounced effect on immobilization of trace metals as trace metals bind with surface functional groups of organic matter, making stable complexes. However, some of the treatments (i.e., 2.5% BC + 2.5% CM amendment ratios of SP-PW and FC-PW soil water status) show a reduced %TOC value under a consecutively low amendment ratio. This is usually attributed to the dissolution and removal of organic carbon particles with the soil solution and degradation by microbial activities.

The NO_3^- concentration in SP and SP-PW treatments shows a pronounced reduction compared with that of the FC and FC-PW treatments (Fig. 3). The saturated condition has a significant impact on the loss of NO_3^- as ammonia. The study of Malhi et al. (1990) noted the extreme loss of NO_3^- from soil under flooded conditions. In our study, nearly twofold concentration of NO_3^- was lost by saturated conditions under the SP and SP-PW treatments. Furthermore, the PO_4^{3-} concentration of SP-PW shows a pronounced increment than that in soil incubated under the other three soil water status. PO_4^{3-} -mineralizing bacteria, including genera *Bacillus*, *Pseudomonas*, and *Rhizobium*, play a critical role in the transformation of insoluble PO_4^{3-} minerals into soluble forms (Rodríguez and Fraga 1999; Qian et al. 2019). Stimulation of PO_4^{3-} -mineralizing microbial growth under water logged conditions, followed by water deficient conditions, might be responsible for the elevated plant available PO_4^{3-} concentrations in SP-PW treatments.

The process of imbibition is involved in the absorption of water to seeds and the activation of enzymes essential for seed germination (Evans and Etherington 1990). The lack of plant available water under permanent wilting point (PW) inhibits the germination of seed, thereby acting as the critical factor for germination. However, there is no significant difference observed for the GI of mung bean and tomato seeds under either saturated point (SP) or field capacity (FC) (Fig. 4). Therefore, trace metal stress has no significant influence for the germination of either mung beans or tomato seeds. However, there was a significant difference observed in the SVI of mung beans between SP and FC (Fig. 5). The lower bioavailability of trace metals under water saturated conditions may have contributed to the significantly high SVI under SP than the SVI of mung beans grown under FC.

5 Conclusions

All the amendment ratios significantly reduced the bioavailable trace metal concentrations we tested under all soil moisture regimes, compared with the respective control. The FC-PW treatment with the highest amendment ratio was significantly effective in reducing the bioavailable fraction of Ni (67.6%), while SP-PW and FC-PW with all amendment ratios ranging from 2.5% BC to 0.0–5.0% CM were equally effective against the removal of Mn (up to 92.1%). The treatment of FC-PW amended with 2.5% BC + 1.0% CM reduced the DTPA extractable Co by 96.9%. Also, the SP soil water content under all amendment ratios (i.e., 2.5% BC + 0.0–5.0% CM) completely immobilized the bioavailable concentration of Cr. The maximum amendment ratio in FC-PW soil was favorable for NO_3^- while SP-PW soil with the highest amendment ratio was most effective at increasing the PO_4^{3-} concentration. Moreover, FC-PW with the maximum amendment ratio contained the highest amount of %TOC, and the maximum CEC value was reported from SP soil with the highest amendment ratio. Besides the amendment ratio, SP and FC influenced GI of mung bean and tomato seeds whereas there was no seed germination observed at PW. However, the SVI of mung bean seedlings exhibited a significant difference ($p < 0.001$) between SP and FC soil water status but no significant difference was detected among different amendment ratios. In contrast, no significant difference of SVI of tomato seedlings was observed ($p = 0.258$) at SP and FC, while a significant difference ($p = 0.020$) of SVI was detected among amendment ratios at SP. Overall, FC-PW soil water status effectively reduced the bioavailable concentrations of all four trace metals examined and increased the soil nutrient levels, including NO_3^- and %TOC. Organic amendment ratios ranging from 2.5% BC + 0.0–5.0% CM effectively reduced the bioavailable fraction of all four trace metals while the highest amendment ratio contributed to the highest soil nutrient content. However, for seedling growth, soil water content was the limiting factor relative to the negative impact of toxic trace metals and the positive effect from increased soil nutrient content.

References

- Agegehu G, Bass AM, Nelson PN, Muirhead B, Wright G, Bird MI (2015) Biochar and biochar-compost as soil amendments: effects on peanut yield, soil properties and greenhouse gas emissions in tropical North Queensland, Australia. *Agric Ecosyst Environ* 213: 72–85
- Alexander EB, Coleman RG, Harrison SP, Keeler-Wolfe T (2007) *Serpentine geocology of western North America: geology, soils, and vegetation*. OUP, USA
- Almaroai YA, Usman AR, Ahmad M, Moon DH, Cho J-S, Joo YK, Jeon C, Lee SS, Ok YS (2014) Effects of biochar, cow bone, and eggshell on Pb availability to maize in contaminated soil irrigated with saline water. *Environ Earth Sci* 71(3):1289–1296
- Anderson JM, Ingram J (1993) *Tropical soil biology and fertility*, 2nd ed. A handbook of methods. CAB international, Wallingford
- Angle JS, Baker AJ, Whiting SN, Chaney RL (2003) Soil moisture effects on uptake of metals by *Thlaspi*, *Alyssum*, and *Berkheya*. *Plant Soil* 256(2):325–332
- Antibachi D, Kelepertzis E, Kelepertsis A (2012) Heavy metals in agricultural soils of the Mouriki-Thiva area (Central Greece) and environmental impact implications. *Soil Sediment Contam* 21(4):434–450
- Avudainayagam S, Megharaj M, Owens G, Kookana RS, Chittleborough D, Naidu R (2003) Chemistry of chromium in soils with emphasis on tannery waste sites: a review. *Rev Environ Contam Toxicol* 178: 53–91
- Aziz RA, Rahim SA, Sahid I, Idris WMR (2015) Speciation and availability of heavy metals on serpentinized paddy soil and paddy tissue. *Procedia Soc Behav Sci* 195:1658–1665
- Baugé S, Lavkulich L, Schreier H (2013) Phosphorus and trace metals in serpentine-affected soils of the Sumas Basin, British Columbia. *Can J Soil Sci* 93(3):359–367
- Baumeister JL, Hausrath EM, Olsen AA, Tschauer O, Adcock CT, Metcalf RV (2015) Biogeochemical weathering of serpentinites: an examination of incipient dissolution affecting serpentine soil formation. *Appl Geochem* 54:74–84
- Beesley L, Moreno-Jiménez E, Gomez-Eyles JL (2010) Effects of biochar and greenwaste compost amendments on mobility, bioavailability and toxicity of inorganic and organic contaminants in a multi-element polluted soil. *Environ Pollut* 158(6):2282–2287
- Beesley L, Inneh OS, Norton GJ, Moreno-Jimenez E, Pardo T, Clemente R, Dawson JJ (2014) Assessing the influence of compost and biochar amendments on the mobility and toxicity of metals and arsenic in a naturally contaminated mine soil. *Environ Pollut* 186:195–202
- Bernal M, Clemente R, Walker D (2007) The role of organic amendments in the bioremediation of heavy metal-polluted soils. In: Gore RW (ed) *Environmental research at the leading edge*. Nova Science Publishers, Inc, New York
- Brooks RR (1987) *Serpentine and its vegetation: a multidisciplinary approach*. Dioscorides Press, Portland
- Cerdeira-Pérez A, Monterroso C, Rodríguez-Garrido B, Machinet G, Echevarria G, Prieto-Fernández Á, Kidd PS (2019) Implementing nickel phytomining in a serpentine quarry in NW Spain. *J Geochem Explor* 197:1–13
- Cheng C-H, Jien S-H, Iizuka Y, Tsai H, Chang Y-H, Hseu Z-Y (2011) Pedogenic chromium and nickel partitioning in serpentine soils along a toposequence. *Soil Sci Soc Am J* 75(2):659–668
- Clemente R, Escolar Á, Bernal MP (2006) Heavy metals fractionation and organic matter mineralisation in contaminated calcareous soil amended with organic materials. *Bioresour Technol* 97(15):1894–1901
- Du Laing G, Vanthuyne D, Vandecasteele B, Tack F, Verloo M (2007) Influence of hydrological regime on pore water metal concentrations in a contaminated sediment-derived soil. *Environ Pollut* 147(3): 615–625
- Evans CE, Etherington JR (1990) The effect of soil water potential on seed germination of some British plants. *New Phytol* 115(3):539–548
- Fengxiang H, Banin A (1997) Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils incubated: I. Under saturated conditions. *Water Air Soil Pollut* 95(1–4):399–423
- Fernandez S, Seoane S, Merino A (1999) Plant heavy metal concentrations and soil biological properties in agricultural serpentine soils. *Commun Soil Sci Plant Anal* 30(13–14):1867–1884
- Galey M, Van Der Ent A, Iqbal M, Rajakaruna N (2017) Ultramafic geocology of south and Southeast Asia. *Bot Stud* 58(1):18
- Gambrell R (1994) Trace and toxic metals in wetlands—a review. *J Environ Qual* 23(5):883–891

- Gray CW, McLaren RG (2006) Soil factors affecting heavy metal solubility in some New Zealand soils. *Water Air Soil Pollut* 175(1–4):3–14
- Gunarathne V, Mayakaduwa S, Vithanage M (2017) Biochar's influence as a soil amendment for essential plant nutrient uptake. In *Essential Plant Nutrients*, pp. 47–67. Springer, Cham
- Han F, Banin A (1997) Long-term transformations and redistribution of potentially toxic heavy metals in arid-zone soils incubated: I. Under saturated conditions. *Water Air Soil Pollut* 95(1):399–423
- Han F, Banin A (2000) Long-term transformations of Cd, Co, Cu, Ni, Zn, V, Mn and Fe in the native arid-zone soils under saturated condition. *Commun Soil Sci Plant Anal* 31:943–957
- Han F, Banin A, Triplett G (2001) Redistribution of heavy metals in arid-zone soils under a wetting-drying cycle soil moisture regime. *Soil Sci* 166(1):18–28
- Han F, Banin A, Kingery W, Triplett G, Zhou L, Zheng S, Ding W (2003) New approach to studies of heavy metal redistribution in soil. *Adv Environ Res* 8(1):113–120
- Harrison SP, Kruckeberg AR (2008) Garden on the rocks. *Nat Hist* 117(4):40–44
- Harrison S, Rajakaruna N (2011) *Serpentine: the evolution and ecology of a model system*. University of California Press, Berkeley, CA
- Herath I, Kumarathilaka P, Navaratne A, Rajakaruna N, Vithanage M (2015) Immobilization and phytotoxicity reduction of heavy metals in serpentine soil using biochar. *J Soils Sediments* 15(1):126–138
- Huang B, Li Z, Huang J, Guo L, Nie X, Wang Y, Zhang Y, Zeng G (2014) Adsorption characteristics of Cu and Zn onto various size fractions of aggregates from red paddy soil. *J Hazard Mater* 264:176–183
- Inyang MI, Gao B, Yao Y, Xue Y, Zimmerman A, Mosa A, Pullammanappallil P, Ok YS, Cao X (2016) A review of biochar as a low-cost adsorbent for aqueous heavy metal removal. *Crit Rev Environ Sci Technol* 46(4):406–433
- Kabala C, Singh BR (2001) Fractionation and mobility of copper, lead, and zinc in soil profiles in the vicinity of a copper smelter. *J Environ Qual* 30(2):485–492
- Kalbitz K, Wennrich R (1998) Mobilization of heavy metals and arsenic in polluted wetland soils and its dependence on dissolved organic matter. *Sci Total Environ* 209(1):27–39
- Kanellopoulos C, Argyraki A, Mitropoulos P (2015) Geochemistry of serpentine agricultural soil and associated groundwater chemistry and vegetation in the area of Atalanti, Greece. *J Geochem Explor* 158:22–33
- Karami N, Clemente R, Moreno-Jiménez E, Lepp NW, Beesley L (2011) Efficiency of green waste compost and biochar soil amendments for reducing lead and copper mobility and uptake to ryegrass. *J Hazard Mater* 191(1–3):41–48
- Kargar M, Clark OG, Hendershot WH, Jutras P, Prasher SO (2015) Immobilization of trace metals in contaminated urban soil amended with compost and biochar. *Water Air Soil Pollut* 226(6):191
- Kashem M, Singh B (2004) Transformations in solid phase species of metals as affected by flooding and organic matter. *Commun Soil Sci Plant Anal* 35(9–10):1435–1456
- Kierczak J, Neel C, Aleksander-Kwaterczak U, Helios-Rybicka E, Bril H, Puziewicz J (2008) Solid speciation and mobility of potentially toxic elements from natural and contaminated soils: a combined approach. *Chemosphere* 73(5):776–784
- Kruckeberg AR (2002) *Geology and plant life: the effects of landforms and rock types on plants*. University of Washington Press, Seattle, WA
- Kumar A, Maiti SK (2013) Availability of chromium, nickel and other associated heavy metals of ultramafic and serpentine soil/rock and in plants. *Int J Adv Res Technol* 3(2):256–268
- Kumarathilaka P, Vithanage M (2017) Influence of *Gliricidia sepium* biochar on attenuate perchlorate-induced heavy metal release in serpentine soil. *J Chem* 2017: 1–8
- Kumarathilaka P, Oze C, Vithanage M (2016) Perchlorate mobilization of metals in serpentine soils. *Appl Geochem* 74:203–209
- Larney FJ, Angers DA (2012) The role of organic amendments in soil reclamation: a review. *Can J Soil Sci* 92(1):19–38
- Lázaro JD, Kidd P, Martínez CM (2006) A phytochemical study of the Trás-os-Montes region (NE Portugal): possible species for plant-based soil remediation technologies. *Sci Total Environ* 354(2–3):265–277
- Liang J, Yang Z, Tang L, Zeng G, Yu M, Li X, Wu H, Qian Y, Li X, Luo Y (2017) Changes in heavy metal mobility and availability from contaminated wetland soil remediated with combined biochar-compost. *Chemosphere* 181:281–288
- Lindsay WL, Norvell WA (1978) Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Sci Soc Am J* 42(3):421–428
- Mackie K, Marhan S, Ditterich F, Schmidt H, Kandeler E (2015) The effects of biochar and compost amendments on copper immobilization and soil microorganisms in a temperate vineyard. *Agric Ecosyst Environ* 201:58–69
- Malhi S, McGill W, Nyborg M (1990) Nitrate losses in soils: effect of temperature, moisture and substrate concentration. *Soil Biol Biochem* 22(6):733–737
- Manios T, Stentiford E, Millner P (2003) Removal of heavy metals from a metaliferous water solution by *Typha latifolia* plants and sewage sludge compost. *Chemosphere* 53(5):487–494
- Neilson S, Rajakaruna N (2015) Phytoremediation of agricultural soils: using plants to clean metal-contaminated arable land. In: *Phytoremediation: Management of Environmental Contaminants*, pp. 159–168, Eds. A. A. Ansari, S. S. Gill, G. R. Lanza, and L. Newman. Springer International Publishing, Switzerland
- Nkrumah PN, Tisserand R, Chaney RL, Baker AJ, Morel JL, Goudon R, Erskine PD, Echevarria G, Van Der Ent A (2019) The first tropical 'metal farm': some perspectives from field and pot experiments. *J Geochem Explor* 198:114–122
- O'dell R, Silk W, Green P, Claassen V (2007) Compost amendment of Cu–Zn minespoil reduces toxic bioavailable heavy metal concentrations and promotes establishment and biomass production of *Bromus carinatus* (Hook and Arn.). *Environ Pollut* 148(1):115–124
- Peijnenburg WJ, Zablotzkaja M, Vijver MG (2007) Monitoring metals in terrestrial environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicol Environ Saf* 67(2):163–179
- Proctor J, Woodell SR (1975) The ecology of serpentine soils. *Adv Ecol Res* 9: 255–366
- Qian T, Yang Q, Jun DCF, Dong F, Zhou Y (2019) Transformation of phosphorus in sewage sludge biochar mediated by a phosphate-solubilizing microorganism. *Chem Eng J* 359:1573–1580
- Rajakaruna N, Baker AJ (2004) Serpentine: a model habitat for botanical research in Sri Lanka. *Ceylon J Sci* 32:1–19
- Rajakaruna N, Bohm BA (2002) Serpentine and its vegetation: a preliminary study from Sri Lanka. *J Appl Bot* 76(1/2):20–28
- Rajakaruna N, Boyd RS (2014) Serpentine soils. In: Gibson D (ed) *Oxford bibliographies in ecology*. Oxford University Press, New York
- Rajapaksha AU, Vithanage M, Oze C, Bandara W, Weerasooriya R (2012) Nickel and manganese release in serpentine soil from the Ussangoda Ultramafic Complex, Sri Lanka. *Geoderma* 189:1–9
- Rizwan M, Ali S, Abbas F, Adrees M, Zia-Ur-Rehman M, Farid M, Gill RA, Ali B (2017) Role of organic and inorganic amendments in alleviating heavy metal stress in oil seed crops. In: *Oil seed crops: yield and adaptations under environmental stress*, pp 224–235, Eds. P. Ahmad. John Wiley & Sons, Ltd
- Rodríguez H, Fraga R (1999) Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol Adv* 17(4–5):319–339
- Schedlbauer JL (2015) Serpentine ecosystem responses to varying water availability and prescribed fire in the US Mid-Atlantic region. *Ecosphere* 6(7):1–14

- Schulz H, Dunst G, Glaser B (2013) Positive effects of composted biochar on plant growth and soil fertility. *Agron Sustain Dev* 33(4):817–827
- Seneviratne M, Seneviratne G, Madawala H, Iqbal M, Rajakaruna N, Bandara T, Vithanage M (2015) A preliminary study of the role of bacterial–fungal co-inoculation on heavy metal phytotoxicity in serpentine soil. *Aust J Bot* 63(4):261–268
- Siddiqui MH, Al-Whaibi MH (2014) Role of nano-SiO₂ in germination of tomato (*Lycopersicon esculentum* seeds Mill.). *Saudi J Biol Sci* 21(1):13–17
- Tang W-W, Zeng G-M, Gong J-L, Liu Y, Wang X-Y, Liu Y-Y, Liu Z-F, Chen L, Zhang X-R, Tu D-Z (2012) Simultaneous adsorption of atrazine and Cu (II) from wastewater by magnetic multi-walled carbon nanotube. *Chem Eng J* 211:470–478
- Tashakor M, Yaacob WZW, Mohamad H (2011) Speciation and availability of Cr, Ni and Co in serpentine soils of Ranau, Sabah. *Am J Geo Sci* 2(1):4–9
- Tashakor M, Hochwimmer B, Brearley FQ (2017) Geochemical assessment of metal transfer from rock and soil to water in serpentine areas of Sabah (Malaysia). *Environ Earth Sci* 76(7):281
- Van Den Berg G, Loch J (2000) Decalcification of soils subject to periodic waterlogging. *Eur J Soil Sci* 51(1):27–33
- Van Gestel CA (2008) Physico-chemical and biological parameters determine metal bioavailability in soils. *Sci Total Environ* 406(3):385–395
- Vashisth A, Nagarajan S (2010) Effect on germination and early growth characteristics in sunflower (*Helianthus annuus*) seeds exposed to static magnetic field. *J Plant Physiol* 167(2):149–156
- Vithanage M, Rajapaksha AU, Oze C, Rajakaruna N, Dissanayake C (2014) Metal release from serpentine soils in Sri Lanka. *Environ Monit Assess* 186(6):3415–3429
- Wei W, Ma R, Sun Z, Zhou A, Bu J, Long X, Liu Y (2018) Effects of mining activities on the release of heavy metals (HMs) in a typical mountain headwater region, the Qinghai-Tibet Plateau in China. *Int J Environ Res Public Health* 15(9):1987
- Wu H, Zeng G, Liang J, Chen J, Xu J, Dai J, Li X, Chen M, Xu P, Zhou Y (2016) Responses of bacterial community and functional marker genes of nitrogen cycling to biochar, compost and combined amendments in soil. *Appl Microbiol Biotechnol* 100(19):8583–8591
- Wu H, Lai C, Zeng G, Liang J, Chen J, Xu J, Dai J, Li X, Liu J, Chen M (2017) The interactions of composting and biochar and their implications for soil amendment and pollution remediation: a review. *Crit Rev Biotechnol* 37(6):754–764
- Xu X, Cao X, Zhao L, Zhou H, Luo Q (2014) Interaction of organic and inorganic fractions of biochar with Pb (II) ion: further elucidation of mechanisms for Pb (II) removal by biochar. *RSC Adv* 4(85):44930–44937
- Ye-Tao T, Teng-Hao-Bo D, Qi-Hang W, Shi-Zhong W, Rong-Liang Q, Ze-Bin W, Xiao-Fang G, Qi-Tang W, Mei L, Tong-Bin C (2012) Designing cropping systems for metal-contaminated sites: a review. *Pedosphere* 22(4):470–488
- Zeng F, Ali S, Zhang H, Ouyang Y, Qiu B, Wu F, Zhang G (2011) The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants. *Environ Pollut* 159(1):84–91
- Zeng G, Wu H, Liang J, Guo S, Huang L, Xu P, Liu Y, Yuan Y, He X, He Y (2015) Efficiency of biochar and compost (or composting) combined amendments for reducing Cd, Cu, Zn and Pb bioavailability, mobility and ecological risk in wetland soil. *RSC Adv* 5(44):34541–34548
- Zhang X, Wang H, He L, Lu K, Sarmah A, Li J, Bolan NS, Pei J, Huang H (2013) Using biochar for remediation of soils contaminated with heavy metals and organic pollutants. *Environ Sci Pollut Res Int* 20(12):8472–8483
- Zheng S, Zhang M (2011) Effect of moisture regime on the redistribution of heavy metals in paddy soil. *J Environ Sci* 23(3):434–443

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.